arXiv:1207.5585v1 [cond-mat.mes-hall] 24 Jul 2012

Sequential Landau-Zener Transitions in A Spin-Phonon Coupled System

Xiaolong Zhang[1,2], Jiahao Huang[1], Yuexia Zhang[2], Kelin Gao[3], and Chaohong Lee[1*]
[1]*State Key Laboratory of Optoelectronic Materials and Technologies,*
School of Physics and Engineering, Sun Yat-Sen University, Guangzhou 510275, China
[2]*Department of Physics, Chongqing University, Chongqing 400044, China and*
[3]*State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics,*
Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China
(Dated: July 25, 2012)

We investigate the Landau-Zener process of a harmonically trapped spin-$\frac{1}{2}$ particle with spin-phonon coupling. Because of the spin-phonon coupling, the energy level structures are modified by the Franck-Condon effects, in which some avoided energy-level-crossings are almost closed and some energy-level-crossings are opened. The close of avoided energy-level-crossings and the open of energy-level-crossings result in the Franck-Condon blockade and the vibrational transitions, respectively. If the spin-phonon coupling is sufficiently strong, the sequential Landau-Zener transitions of ladder-like population transition appear in the Landau-Zener process, in which an initially ground-state system is driven from one far-off-resonance limit to the other far-off-resonance limit. We derive a generalized Landau-Zener formula which is well consistent with our numerical results.

PACS numbers: 03.75.Lm, 33.70.Ca, 33.20.Wr, 03.65.Xp

Landau-Zener (LZ) problem [1, 2], a well-known fundamental problem in time-dependent quantum mechanics, concerns how non-adiabatic transition appears in a two-level system driven through an avoided energy-level-crossing. According to the quantum adiabatic theorem [3, 4], if the system varies infinitely slow, non-adiabatic transition will not take place and the system will always be in an eigenstate of its instantaneous Hamiltonian. The studies of LZ transition are not only of great fundamental interests [5–9], but also of extensive applications [10–17] in quantum state engineering, quantum interferometry and quantum computation etc.

Up to now, there are lots of theoretical and experimental studies of LZ transitions in systems of decoupled internal spin states and external motional states. However, to the best of our knowledge, there are almost no studies on LZ transitions in closed quantum systems whose internal spin states and external motional states are coupled. Spin-phonon coupling (SPC) is a typical coupling between internal spin states and external motional states. *How SPC affects a LZ process? What does SPC bring to a conventional LZ transition?*

It has been demonstrated that strong SPC may induce Franck-Condon (FC) blockade and vibrational sidebands. The FC principle [18, 19] tells us that the FC blockade takes place if the FC factor, which is defined as the square of the overlap integral between the vibrational wave-functions of the two involved states, is sufficiently small to be ignorable. The FC blockade induced by strong SPC has been found in several systems, such as molecular junctions [20–22], nano-tube quantum dots [23–26], neutral atoms [27], and a single trapped ion [28]. On the other hand, nonzero FC factors may cause vibrational sidebands [18, 19, 23, 27, 29–32], that is, the population transfer or the electronic tunnelling has been shown to excite vibrational modes. *Is there any signature of the FC effects in the LZ process of a spin-phonon coupled system?*

In this Letter, we investigate the LZ process of a spin-phonon coupled system of a single spin-$\frac{1}{2}$ particle in a spin-dependent harmonic potential. We explore how SPC affects energy-level structures and the LZ process. The energy gaps of avoided energy-level-crossings become narrow when the SPC becomes strong. At the same time, some energy-level-crossings are gradually opened and then closed. The appearance of FC blockade and vibrational sidebands are direct results of the close of avoided energy-level-crossings and the open of energy-level-crossings, respectively. Under sufficiently strong SPCs, in contrast to the LZ transition in a system without SPC, the sequential LZ transitions of ladder-like population transition appear. We find that the FC blockade corresponds to the absence of some specific population steps. By treating the sequential LZ transitions as a sequence of conventional two-level LZ transitions and applying the conventional two-level LZ formula again and again, we obtain a generalized LZ formula for the final populations. Based upon the experimental techniques for exploring the FC blockade, it is possible to test our prediction of the sequential LZ transitions in future experiments.

We consider a harmonically trapped particle, which may be an electron, an atom, a molecular or a quantum dot and only two of its internal spin states of different magnetic dipole moments are involved. Below, we call this particle as a spin-$\frac{1}{2}$ particle and denote two involved spin states as spin-up state $|\uparrow\rangle$ and spin-down state $|\downarrow\rangle$. In the LZ process, the two spin states are coupled by lasers with a linearly sweeping detuning. For simplicity and without loss of generality, we concentrate our studies on one-dimension systems with SPCs induced by gradient Zeeman fields. The Hamiltonian for our spin-phonon coupled system reads as,

$$H = H_{LZ} + H_{ho} + H_{spc}, \tag{1}$$

where $H_{LZ} = -\frac{\hbar\delta(t)}{2}\sigma_z - \frac{\hbar\Omega}{2}\sigma_x$ with $\delta(t) = \delta_0 + \alpha t$ is the conventional LZ Hamiltonian, $H_{ho} = -\frac{\hbar^2}{2m}\frac{\partial^2}{\partial z^2} + \frac{1}{2}m\omega_z^2 z^2$ describes the external motion and $H_{spc} = m\omega_z^2 z_0 z \sigma_z$ characterizes the SPC. Here, $\sigma_{x,z}$ are Pauli matrices, $\hbar$ is the Planck constant, m is the particle mass, Ω is the Rabi frequency, $\delta(t)$ is the detuning, and z_0 is the SPC strength. Taking into account the SPC, the particle feels a spin-dependent potential $V_\sigma(z,t) = -\frac{1}{2}m\omega_z^2 z_0^2 + \frac{1}{2}m\omega_z^2\left(z + \langle\sigma_z\rangle z_0\right)^2 - \frac{\hbar\delta(t)}{2}\langle\sigma_z\rangle$ with $\langle\sigma_z\rangle = +1$ for $\sigma =\uparrow$ and $\langle\sigma_z\rangle = -1$ for $\sigma =\downarrow$. In Fig. 1, we show the schematic diagram for the LZ process of Hamiltonian (1).



FIG. 1: Schematic diagram for the Landau-Zener process of a spin-$\frac{1}{2}$ particle in a harmonic trap. The red dashed curves and the blue solid curves denote the spin-dependent harmonic trap $V_\uparrow(z)$ and $V_\downarrow(z)$, respectively. The detuning is linearly swept from one far-off-resonance limit (left column) to the other far-off-resonance limit (right column) through the zero point (middle column). (a) $z_0 = 0$ for the case of no spin-phonon coupling and (b) $z_0 \neq 0$ for the case of spin-phonon coupling.

Denoting the quantum state of Hamiltonian (1) as $|\Psi\rangle = \begin{pmatrix}\psi_\uparrow(z,t)\\ \psi_\downarrow(z,t)\end{pmatrix}$, its time-evolution obeys the coupled Schrödinger equations,

$$i\hbar\frac{\partial}{\partial t}\psi_\uparrow = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial z^2} + V_\uparrow(z,t)\right]\psi_\uparrow - \frac{\hbar\Omega}{2}\psi_\downarrow, \tag{2}$$

$$i\hbar\frac{\partial}{\partial t}\psi_\downarrow = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial z^2} + V_\downarrow(z,t)\right]\psi_\downarrow - \frac{\hbar\Omega}{2}\psi_\uparrow, \tag{3}$$

with $\int dz\left(|\psi_\uparrow(z,t)|^2 + |\psi_\downarrow(z,t)|^2\right) = 1$ imposed by the normalization condition. Given the eigenstates $|\sigma n\rangle$ for $V_\sigma(z,t)$, the wave-functions can be expressed as $\psi_\sigma(z,t) = \sum_{n=0}^{+\infty} C_{\sigma n}(t)|\sigma n\rangle$ with $C_{\sigma n}(t) = \langle n\sigma|\psi_\sigma(z,t)\rangle$. In the basis composed of $|\sigma n\rangle$, the system obeys an effective Hamiltonian,

$$H = \sum_{\sigma,n} E_{\sigma n}|\sigma n\rangle\langle n\sigma| - \frac{\Omega}{2}\sum_{n',n}\sqrt{F_{n'n}}\left[|\downarrow n'\rangle\langle n\uparrow| + h.c.\right]. \tag{4}$$

Here, the FC factors $F_{n'n} = |\langle n'\downarrow|\uparrow n\rangle|^2 = |\langle n'\uparrow|\downarrow n\rangle|^2$ and $E_{\sigma n} = n\hbar\omega_z - \langle\sigma_z\rangle\hbar\delta/2$ with $\langle\sigma_z\rangle = +1$ for $\sigma =\uparrow$ and $\langle\sigma_z\rangle = -1$ for $\sigma =\downarrow$. Comparing with Hamiltonian (1), the zero-energy point of Hamiltonian (4) is shifted to $\left(\hbar\omega_z - m\omega_z^2 z_0^2\right)/2$. From the effective Hamiltonian (4), the complex amplitudes $C_{\sigma n}(t)$ obey

$$i\hbar\frac{\partial}{\partial t}C_{\uparrow n}(t) = E_{\uparrow n}C_{\uparrow n}(t) - \frac{\hbar\Omega}{2}\sum_{n'}\sqrt{F_{n'n}}C_{\downarrow n}(t), \tag{5}$$

$$i\hbar\frac{\partial}{\partial t}C_{\downarrow n}(t) = E_{\downarrow n}C_{\downarrow n}(t) - \frac{\hbar\Omega}{2}\sum_{n'}\sqrt{F_{n'n}}C_{\uparrow n}(t). \tag{6}$$

Obviously, if $F_{n'n} = 0$, there will be no population transfer between $|\sigma n\rangle$ and $|\sigma' n'\rangle$ with $\sigma \neq \sigma'$.

Below, we analyze the time evolutions and the population transitions in the LZ process. By numerically integrating the coupled Schrödinger equations (2) and (3), we obtain $\psi_\sigma(z,t)$ and then calculate both the spin populations $P_\sigma(t) = \int dz\,|\psi_\sigma(z,t)|^2$ and the phonon populations $P_{\sigma n}(t) = |C_{\sigma n}(t)|^2$. In our numerical simulation, we have chosen the natural units of $m = 1$, $\hbar = 1$ and $\omega_z = 1$. The initial state is chosen as the ground state in the negative far-off-resonance limit ($\delta < 0$ and $|\delta| \gg \Omega$), that is, $\psi_\downarrow(z, t = 0) = |\downarrow 0\rangle$ and $\psi_\uparrow(z, t = 0) = 0$. In the LZ processes, the Rabi frequency Ω is fixed as 0.2, the detuning is linearly swept according to $\delta = \delta_0 + \alpha t$ with the initial detuning $\delta_0 = -10$ and the sweeping rate $\alpha = 2.5 \times 10^{-4}$.



FIG. 2: Time evolution of probability densities in the Landau-Zener process. The left column is $|\psi_\downarrow(z,t)|^2$ and the right column is $|\psi_\uparrow(z,t)|^2$. The top, middle and bottom rows correspond to the SPC strength $z_0 = 0$, 1.5 and 2.5, respectively. The other physical parameters are chosen as $\delta_0 = -10$, $\alpha = 2.5 \times 10^{-4}$ and $\Omega = 0.2$.

In Fig. 2, for different values of the SPC strength z_0, we show the time evolution of the probability densities

$|\psi_{\uparrow,\downarrow}(z,t)|^2$ in the LZ process. If there is no SPC, the system undergoes adiabatic evolution and the two probability densities keep in Gaussian shapes. The unchanged shapes of $|\psi_{\uparrow,\downarrow}(z,t)|^2$ for $z_0 = 0$ indicate the absence of phonon excitations. When the SPC strength z_0 increases, although $|\psi_{\downarrow}(z,t)|^2$ keeps in a Gaussian shape, multi-hump structures gradually appear in $|\psi_{\uparrow}(z,t)|^2$. The appearance of multi-hump structures in $|\psi_{\uparrow}(z,t)|^2$ is a signature of phonon excitations. In particular, the significant changes of the two probability densities sequentially take place in the vicinity of $\delta = n\hbar\omega$ (where n are non-negative integers).



FIG. 3: Sequential Landau-Zener transitions induced by spin-phonon coupling. (a) The populations $P_{\downarrow}$ (red curves) and $P_{\uparrow}$ (blue curves) versus the detuning δ for $z_0 = 0$ and (b) the ones for $z_0 = 2.5$. (c) and (d) are the two energy spectra corresponding to (a) and (b), respectively. The sequential Landau-Zener transitions in (b) have ladder-like population transition with steps at $\delta = n\hbar\omega_z$. The numbers 1 to 7 in (d) label the avoided energy-level-crossings, which correspond to the population steps in (b).

In Fig. 3 (a), we show the population transitions corresponding to the time evolutions in the top row of Fig. 2. Due to the absence of SPC, the phonon states are spin-independent and the FC factors $F_{n',n}$ are non-zero if and only if $n' = n$. Therefore, the time evolution of spin states and phonon states are decoupled and phonon excitations will not take place in the LZ process. In the corresponding energy spectrum, the avoided energy-level-crossings only appear around $\delta = 0$ and these avoided energy-level-crossings dominate the population transfer in the LZ process, see Fig. 3 (c). In the LZ process, as labeled by the arrows, the system evolves along its instantaneous ground state due to the sweeping rate $\alpha = 2.5 \times 10^{-4}$ is sufficiently small.

In Fig. 3 (b), we show the population transitions corresponding to the time evolutions in the bottom row of Fig. 2. Due to the strong SPC, the FC blockade and the sequential LZ transitions appear in the LZ process. Corresponding to the significant density changes in Fig. 2, a series of population steps appear at $\delta = n\hbar\omega$. This ladder-like population transition is a direct signature of the sequential LZ transitions. In addition, unlike the conventional LZ transition, there is no population step at $\delta = 0$. The absence of the population step at $\delta = 0$ is a result of the FC blockade between the lowest-phonon states $|\downarrow 0\rangle$ and $|\uparrow 0\rangle$. Based upon our numerical results for different values of z_0 (not shown here), the FC blockade appear only when z_0 is sufficiently large and more population steps will disappear due to the FC blockades between the lowest-phonon state $|\downarrow 0\rangle$ and the high-phonon states $|\uparrow n\rangle$ take place for larger z_0. In the energy spectrum, because of the FC effects, some avoided energy-level-crossings are almost closed due to the corresponding FC factors are very small, and some energy-level-crossings opened due to the corresponding FC factors become non-zero, see Fig. 3 (d). Therefore, as labeled by the arrows, the system undergoes sequential LZ transitions in which sequential phonon excitations accompany the ladder-like spin population transition.

To characterize the phonon excitations, one has to analyze the population dynamics of $|\sigma n\rangle$. Given $\psi_{\sigma}(z,t)$, it is easy to obtain the phonon populations $P_{\sigma n}(t) = |C_{\sigma n}(t)|^2$ with $C_{\sigma n}(t) = \langle n\sigma|\psi_{\sigma}(z,t)\rangle$. In the absence of SPC, the phonon populations $P_n(t) = P_{\downarrow n}(t) + P_{\uparrow n}(t)$ keep unchanged, i.e. $P_n(t) = P_n(0)$. In the presence of SPC, due to the occurrence of the phonon excitations from $|\downarrow 0\rangle$ to $|\uparrow n\rangle$ around $\delta = n\hbar\omega_z$, step-like changes appear in the phonon populations. In other words, the population step at $\delta = n\hbar\omega_z$ is caused by the sideband transition between $|\downarrow 0\rangle$ and $|\uparrow n\rangle$.

The sequential LZ transitions can be treated as a sequence of conventional two-level LZ transitions. To calculate the transition probability of a two-level LZ transition in the sequence, in addition to the sweeping rate α, one has to know the energy gap for its avoided energy-level-crossing. The energy gap sensitively depends on the SPC strength z_0. By diagonalizing the effective Hamiltonian (4), the energy gap for the avoided energy-level-crossing between $|\downarrow n\rangle$ and $|\uparrow n'\rangle$ is given as $\Delta_{n'n} = \Omega\sqrt{F_{n'n}}$. If there is no SPC, $\Delta_{n'n} = \Omega\delta_{n'n}$ with $\delta_{n'n}$ denoting a Kronecker delta function. If the SPC is sufficiently weak, Δ_{00} is still large enough and the LZ process is similar to the conventional LZ process of no SPC. The energy gaps Δ_{nn} become narrow when the SPC becomes strong and at the same time the other energy gaps $\Delta_{n'n}$ of $n' \neq n$ are gradually opened and then closed. If the SPC is sufficiently strong, the almost vanishing $\Delta_{n'n}$ may induce the blockade of the vibrational transitions $|\downarrow n\rangle \Leftrightarrow |\uparrow n'\rangle$ in the LZ process.

Now, by using the conventional two-level LZ formula again and again, we derive the analytical formula for the final populations. The sequential LZ transitions can be decomposed into a sequence of conventional two-level LZ transitions: $|\downarrow 0\rangle \Leftrightarrow |\uparrow 0\rangle$, $|\downarrow 0\rangle \Leftrightarrow |\uparrow 1\rangle$, $\cdots$,$|\downarrow 0\rangle \Leftrightarrow |\uparrow n\rangle$, $\cdots$, see Fig. 3 (d). Therefore, by applying the conventional two-level LZ formula one by one, the final populations are given as

$$\begin{aligned} P_{\uparrow 0} &= 1 - P_{LZ}(\Delta_{00}, \alpha), \\ P_{\uparrow 1} &= P_{LZ}(\Delta_{00}, \alpha)(1 - P_{LZ}(\Delta_{01}, \alpha)), \\ P_{\uparrow 2} &= P_{LZ}(\Delta_{00}, \alpha) P_{LZ}(\Delta_{01}, \alpha)(1 - P_{LZ}(\Delta_{02}, \alpha)), \\ &\cdots, \\ P_{\uparrow n} &= \left[\prod_{n'=0}^{n'=n-1} P_{LZ}(\Delta_{0n'}, \alpha) \right] (1 - P_{LZ}(\Delta_{0n}, \alpha)), \quad (7) \\ &\cdots, \\ P_{\downarrow n} &= \begin{cases} \prod_{n'=0}^{n'=+\infty} P_{LZ}(\Delta_{0n'}, \alpha), & \text{for} \quad n = 0, \\ 0, & \text{for} \quad n \neq 0, \end{cases} \end{aligned}$$

with $P_{LZ}(\Delta_{0n}, \alpha) = \exp\left(-\frac{\pi \Delta_{0n}^2}{2\alpha}\right)$ given by the conventional two-level LZ formula.

TABLE I: Comparison between numerical and analytical results of final populations. The first column is the SPC strength z_0, the second column is the FC factor, the third column is the final population obtained from numerical integration, and the last column is the final population given by the generalized LZ formula (7). The rows of $z_0 = 0$ and 2.5 correspond to (a) and (b) of Fig. 3, respectively.

SPC	FC Factor	Numerical	Analytical
$z_0 = 0$	$F_{00} = 1$	$P_{\uparrow 0} = 0.9998$	$P_{\uparrow 0} = 1.0000$
$z_0 = 2.5$	$F_{00} = 3.7 \times 10^{-6}$	$P_{\uparrow 0} = 0.0005$	$P_{\uparrow 0} = 0.0005$
	$F_{01} = 4.7 \times 10^{-5}$	$P_{\uparrow 1} = 0.0059$	$P_{\uparrow 1} = 0.0058$
	$F_{02} = 0.0003$	$P_{\uparrow 2} = 0.0367$	$P_{\uparrow 2} = 0.0357$
	$F_{03} = 0.0012$	$P_{\uparrow 3} = 0.1383$	$P_{\uparrow 3} = 0.1355$
	$F_{04} = 0.0038$	$P_{\uparrow 4} = 0.3152$	$P_{\uparrow 4} = 0.3117$
	$F_{05} = 0.0095$	$P_{\uparrow 5} = 0.3535$	$P_{\uparrow 5} = 0.3556$
	$F_{06} = 0.0197$	$P_{\uparrow 6} = 0.1378$	$P_{\uparrow 6} = 0.1423$
	$F_{07} = 0.0353$	$P_{\uparrow 7} = 0.0121$	$P_{\uparrow 7} = 0.0128$

The final populations given by the generalized LZ formula (7) are well consistent with our numerical results. In Table I, for different values of the SPC strength z_0, we compare the final populations estimated by the generalized LZ formula (7) with the corresponding ones obtained by numerical integration. For $z_0 = 0$, which corresponds to the numerical results shown in Fig. 3 (a), the phonon number n is exactly truncated at 0 because there is no SPC. For $z_0 = 2.5$, which corresponds to the numerical results shown in Fig. 3 (b), the phonon number n can be approximately truncated at 7. In particular, the FC factor F_{00} is in order of 10^{-6} and the final population $P_{\uparrow 0}$ is in order of 10^{-4}. As the initial state is $|\downarrow 0\rangle$, such a small $P_{\uparrow 0}$ is a signature of the FC blockade between $|\downarrow 0\rangle$ and $|\uparrow 0\rangle$. For a stronger SPC, there may appear more FC blockades between $|\downarrow 0\rangle$ and $|\uparrow n\rangle$ with n up to a larger integer number. The numerical and analytical data of the final populations clearly show that the largest absolute difference is less than 0.005 and the largest relative difference is less than 5.5%. This means that the generalized LZ formula (7) is a very good estimation for the final populations.

By using a single electron spin in suspended carbon nanotube quantum dot [24], a single ion in spin-dependent Paul trap [28], or ultracold atoms in spin-dependent optical lattices [27], it is possible to test our prediction of the sequential LZ transitions. Here, for two examples, we briefly discuss the experimental possibility via a single trapped ion [28] or noninteracting ultracold atoms [27] with tunable SPCs. In the case of a single trapped ion, the spin-dependent potential can be formed by imposing a gradient magnetic field on the Paul trap [33] and the SPC strength is determined by the field gradient. In the case of noninteracting ultracold atoms, the spin-dependent optical lattices [34] can be created by two polarized counter-propagating laser beams and the SPC strength can be adjusted by tuning the polarization angle. In both two cases, the two internal spin states can be coupled by Raman laser beams and the detuning can be varied by modifying the laser frequency.

In summary, we have studied the LZ problem of a harmonically trapped spin-$\frac{1}{2}$ particle with SPC. The intrinsic mechanism of the sequential LZ transitions and the direct signatures of the FC effects in the LZ process have been explored. Due to the FC effects, the energy gap for the avoided energy-level-crossing between $|\downarrow n\rangle$ and $|\uparrow n'\rangle$ is given as $\Delta_{n'n} = \Omega\sqrt{F_{n'n}}$. The sequential LZ transitions, which have ladder-like population transition, take place when the SPC is sufficiently strong. The presence of population steps at $\delta = n\hbar\omega_z$ is a signature of the vibrational sidebands and the absence of some specific steps is a signature of the FC blockade. The sequential LZ transitions can be treated as a sequence of conventional two-level LZ transitions. We analytically derive a generalized LZ formula, which gives an excellent estimation for the final populations. We also briefly discuss the experimental possibility.

This work is supported by the National Basic Research Program of China under Grant No. 2012CB821300, the National Natural Science Foundation of China under Grants No. 11075223 and No. 10804132, the Program for New Century Excellent Talents in University of Ministry of Education of China under Grant No. NCET-10-0850 and the Fundamental Research Funds for Central Universities of China.

---

* Corresponding author. Email: chleecn@gmail.com

[1] L. Landau, Phys. Z. Sowjetunion **2**, 46 (1932).

[2] C. Zener, Proc. R. Soc. A **137**, 696 (1932).

[3] M. Born and V. A. Fock, Zeitschrift für Physik A **51**, 165 (1928).

[4] T. Kato, J. Phys. Soc. Jpn. **5**, 435 (1950).

[5] P. Kral, I. Thanopulos, and M. Shapiro, Rev. Mod. Phys. **79**, 53 (2007).

[6] S. N. Shevchenko, S. Ashhabb, and F. Nori, Phys. Rep. **492**, 1 (2010).

[7] A. Altland and V. Gurarie, Phys. Rev. Lett. **100**, 063602 (2008).

[8] J. Keeling and V. Gurarie, Phys. Rev. Lett. **101**, 033001 (2008).

[9] Y. Chen, S. D. Huber ,S. Trotzky, I. Bloch, and E. Altman, Nature Phys. **7**, 61 (2010).

[10] Y. Makhlin, G. Schön, and A. Shnirman, Rev. Mod. Phys. **73**, 357 (2001).

[11] W.D. Oliver, Y. Yu, J. C. Lee, K. K. Berggren, L. S. Levitov, T. P. Orlando, Science **310**, 1653 (2005).

[12] A. O. Niskanen, K. Harrabi, F. Yoshihara, Y. Nakamura, S. Lloyd, J. S. Tsai, Science **316**, 723 (2007).

[13] L. F. Wei, J. R. Johansson, L. X. Cen, S. Ashhab, and F. Nori, Phys. Rev. Lett. **100**, 113601 (2008).

[14] M. D. LaHaye, J. Suh, P. M. Echternach, K. C. Schwab, and M. L. Roukes, Nature **459**, 960 (2009).

[15] J. R. Petta, H. Lu, and A. C. Gossard, Science **327**, 669 (2010).

[16] J. Q. You and F. Nori, Nature **474**, 589 (2011).

[17] M. G. Bason, M. Viteau, N. Malossi, P. Huillery, E. Arimondo, D. Ciampini, R. Fazio, V. Giovannetti, R. Mannella, and O. Morsch, Nature Phys. **8**, 147 (2012).

[18] J. Franck, Trans. Farad. Soc. **21** 536 (1926).

[19] E. Condon, Phys. Rev. **28** 1182 (1926).

[20] J. Koch, and F. von Oppen, Phys. Rev. Lett. **94**, 206804 (2005).

[21] J. Koch, F. von Oppen, and A. V. Andreev, Phys. Rev. B **74**, 205438 (2006).

[22] C. Joachim, J. K. Gimzewski, and A. Aviram, Nature **408**, 541 (2000).

[23] R. Leturcq, C. Stampfer, K. Inderbitzin, L. Durrer, C. Hierold, E. Mariani, M. G. Schultz, F. von Oppen and K. Ensslin, Nature Phys. **5**, 327 (2009).

[24] A. Pályi, P. R. Struck, M. Rudner, K. Flensberg, and G. Burkard, Phys. Rev. Lett. **108**, 206811 (2012).

[25] C. Ohm, C. Stampfer, J. Splettstoesser, and M. R. Wegewijs, Appl. Phys. Lett. **100**, 143103 (2012).

[26] K. L. Ekinci, and M. L. Roukes, Rev. Sci. Instrum. **76**, 061101 (2005)

[27] L. Förster, M. Karski, J.-M. Choi, A. Steffen, W. Alt, D. Meschede, A. Widera, E. Montano, J. H. Lee, W. Rakreungdet, and P. S. Jessen, Phys. Rev. Lett. **103**, 233001 (2009).

[28] Y. M. Hu, W. L. Yang, Y. Y. Xu, F. Zhou, L. Chen, K. L. Gao, M. Feng, and C. Lee, New J. Phys. **13**, 053037 (2011).

[29] H. Park, J. Park, A. K. L. Lim, E. H. Anderson, A. P. Alivisatos, and P. L. McEuen, Nature **407**, 57 (2000).

[30] L. H. Yu, Z. K. Keane, J. W. Ciszek, L. Cheng, M. P. Stewart, J. M. Tour, and D. Natelson, Phys. Rev. Lett. **93**, 266802 (2004).

[31] A. N. Pasupathy, J. Park, C. Chang, A. V. Soldatov, S. Lebedkin, R. C. Bialczak, J. E. Grose, L. A. K. Donev, J. P. Sethna, D. C. Ralph, and P. L. McEuen, Nano Lett. **5**, 203 (2005).

[32] S. Sapmaz, P. Jarillo-Herrero, Ya. M. Blanter, C. Dekker, and H. S. J. van der Zant, Phys. Rev. Lett. 96, 026801 (2006).

[33] F. Mintert and C. Wunderlich, Phys. Rev. Lett. 87, 257904 (2001).

[34] O. Mandel, M. Greiner, A. Widera, T. Rom, T. W. Hänsch, and I. Bloch,Phys. Rev. Lett. **91**, 010407 (2003).